

Shilla distance-regular graphs

Jack H. Koolen, Jongyook Park

Department of Mathematics, POSTECH, Pohang, 790-784, Korea

E-mail: koolen@postech.ac.kr, jongyook@postech.ac.kr

Abstract

A Shilla distance-regular graph Γ (say with valency k) is a distance-regular graph with diameter 3 such that its second largest eigenvalue equals to a_3 . We will show that a_3 divides k for a Shilla distance-regular graph Γ , and for Γ we define $b = b(\Gamma) := \frac{k}{a_3}$. In this paper we will show that there are finitely many Shilla distance-regular graphs Γ with fixed $b(\Gamma) \geq 2$. Also, we will classify Shilla distance-regular graphs with $b(\Gamma) = 2$ and $b(\Gamma) = 3$. Furthermore, we will give a new existence condition for distance-regular graphs, in general.

Key Words: distance-regular graph; Existence condition; Terwilliger graph

2000 Mathematics Subject Classification: 05E30

1 Introduction

In this paper we study distance-regular graphs Γ with diameter 3. (For definitions, see next section.) For a distance-regular graph with diameter 3, we will show that the second largest eigenvalue θ_1 is at least $\min\{\frac{a_1 + \sqrt{a_1^2 + 4k}}{2}, a_3\}$, where k is the valency (see, Lemma 6 below), and that $\theta_1 = a_3$ if and only if $\theta_1 = \frac{a_1 + \sqrt{a_1^2 + 4k}}{2}$. A distance-regular graph Γ with diameter 3 is called *Shilla* if $\theta_1 = a_3$. It follows that for a Shilla distance-regular graph Γ , a_3 divides k and we will put $b(\Gamma) := \frac{k}{a_3}$. In this paper we will show that there exist finitely many (non-isomorphic) Shilla distance-regular graphs with fixed $b(\Gamma) \geq 2$. This result relies on a new existence condition, Theorem 4, for distance-regular graphs. Furthermore we will classify Shilla distance-regular graphs Γ with $b(\Gamma) \in \{2, 3\}$.

This paper is organized as follows: In Section 2, we will give definitions. In Section 3, we give the new existence condition for distance-regular graphs, and in Section 4 we will discuss Shilla distance-regular graphs.

2 Definitions and preliminaries

Suppose that Γ is a connected graph with the vertex set $V(\Gamma)$ and the edge set $E(\Gamma)$, where $E(\Gamma)$ consists of the unordered pairs of adjacent two vertices. The distance $d_\Gamma(x, y)$ between any two vertices x, y of Γ is the length of a shortest path between x and y in Γ .

Let Γ be a connected graph. For a vertex $x \in V(\Gamma)$, define $\Gamma_i(x)$ to be the set of vertices which are at distance precisely i from x ($0 \leq i \leq D$) where $D := \max\{d_\Gamma(x, y) \mid x, y \in V(\Gamma)\}$ is the diameter of Γ . In addition, define $\Gamma_{-1}(x) := \emptyset$ and $\Gamma_{D+1}(x) := \emptyset$. We will write $\Gamma(x)$ instead of $\Gamma_1(x)$ and we denote $x \sim_\Gamma y$ or simply $x \sim y$ if two vertices x and y are adjacent in Γ . For $x_1, x_2, \dots, x_l \in V(\Gamma)$, define

$$\Gamma(x_1, x_2, \dots, x_l) := \bigcap_{i=1}^l \Gamma(x_i).$$

A connected graph Γ with diameter D is called *distance-regular* if there are integers b_i, c_i ($0 \leq i \leq D$) such that for any two vertices $x, y \in V(\Gamma)$ with $d_\Gamma(x, y) = i$, there are precisely c_i neighbors of y in $\Gamma_{i-1}(x)$ and b_i neighbors of y in $\Gamma_{i+1}(x)$. In particular, distance-regular graph Γ is regular with valency $k := b_0$ and we define $a_i := k - b_i - c_i$ for notational convenience. Note that $a_i = |\Gamma(y) \cap \Gamma_i(x)|$ holds for any two vertices x, y with $d_\Gamma(x, y) = i$ ($0 \leq i \leq D$). For a distance-regular graph Γ and a vertex $x \in V(\Gamma)$, we denote $k_i := |\Gamma_i(x)|$. The numbers a_i, b_{i-1} and c_i ($1 \leq i \leq D$) are called the *intersection numbers* of Γ , and they satisfy the following three conditions:

$$(i) \quad k = b_0 > b_1 \geq \dots \geq b_{D-1};$$

$$(ii) \quad 1 = c_1 \leq c_2 \leq \dots \leq c_D;$$

$$(iii) \quad b_i \geq c_j \text{ if } i + j \leq D.$$

The array $\{b_0, b_1, \dots, b_{D-1}; c_1, c_2, \dots, c_D\}$ is called the *intersection array* of a distance-regular graph Γ . Suppose that Γ is a distance-regular graph with valency $k \geq 2$ and diameter $D \geq 2$, and let A_i be the matrix of Γ such that the rows and the columns of A_i are indexed by $V(\Gamma)$ and the (x, y) -entry of A_i equals 1 whenever $d_\Gamma(x, y) = i$ and 0 otherwise. We will denote the adjacency matrix of Γ as A instead of A_1 . Then Γ has exactly $(D + 1)$ distinct eigenvalues, say $k = \theta_0 > \theta_1 > \dots > \theta_D$, and let m_i be the multiplicity of θ_i ($0 \leq i \leq D$), where an eigenvalue of Γ is that of A .

For an eigenvalue θ of Γ , the sequence $u_0 = u_0(\theta) = 1, u_1 = u_1(\theta) = \frac{\theta}{k}, u_i = u_i(\theta)$ ($2 \leq i \leq D$) satisfying

$$c_i u_{i-1}(\theta) + a_i u_i(\theta) + b_i u_{i+1}(\theta) = \theta u_i(\theta)$$

is called the *standard sequence* corresponding to the eigenvalue θ .

N. Biggs[1, p.131] showed that for an eigenvalue θ of a distance-regular graph Γ , its multiplicity m is given by

$$m = \frac{|V(\Gamma)|}{\sum_{i=0}^D k_i u_i(\theta)^2}. \quad (1)$$

The Bose-Mesner algebra M for a distance-regular graph Γ is the matrix algebra generated by the adjacency matrix A of Γ . A basis of M is $\{A_i \mid i = 0, \dots, D\}$, where $A_0 = I$. The algebra M has also a basis consisting of primitive idempotents $\{E_0 = \frac{1}{n}J, E_1, \dots, E_D\}$, where $n = |V(\Gamma)|$ and E_i is the orthogonal projection onto

the eigenspace of θ_i . Under the componentwise multiplication \circ , $E_i \circ E_j = \frac{1}{n} \sum_{k=0}^D q_{ij}^k E_k$.

The numbers q_{ij}^k ($0 \leq i, j, k \leq D$) are called the *Krein parameters* of Γ and are always non-negative by Delsarte [1, Theorem 2.3.2]. We say that Γ is *Q-polynomial* if there is an order of the primitive idempotents $E_0 = \frac{1}{n}J, E_1, \dots, E_D$ such that $q_{1j}^k = 0$ if $|j - k| > 1$. We say that Γ is *Q-polynomial* with respect to θ if E_1 is the orthogonal projection on the eigenspace of θ .

In this paper we say that an intersection array is *feasible* if it satisfies the following four conditions:

- (i) all its intersection numbers p_{jl}^i are integral;
(where $p_{jl}^i = |\{z \mid d_\Gamma(x, z) = j, d_\Gamma(y, z) = l\}|$ for any vertices x and y at distance i)
- (ii) all the multiplicities are positive integers;
- (iii) for any $0 \leq i \leq D$, $k_i a_i$ is even;
- (iv) all Krein parameters are non-negative.

Recall that a *clique* of a graph is a set of mutually adjacent vertices and that a *co-clique* of a graph is a set of vertices with no edges. For a graph Γ , the *local graph* at a vertex $x \in V(\Gamma)$ is the subgraph induced by $\Gamma(x)$ in Γ and we denote it by $\Delta(x)$. Let Δ be a graph. We say Γ is locally Δ if the local graph $\Delta(x)$ is isomorphic to Δ for all vertices $x \in V(\Gamma)$. An *order(s, t)-graph* is a graph such that each $\Delta(x)$ is the disjoint union of $t + 1$ copies of $(s + 1)$ -cliques. A *Terwilliger graph* is a connected non-complete graph Γ such that, for any two vertices u, v at distance two, the subgraph induced by $\Gamma(u, v)$ in Γ is a clique of size μ (for some fixed $\mu \geq 1$).

Recall the following interlacing result.

Theorem 1 (cf. Haemers[3]) Let A be a real symmetric $n \times n$ matrix and let B be a principal submatrix of A with order $m \times m$. Then, for $i = 1, \dots, m$,

$$\theta_{n-m+i}(A) \leq \theta_i(B) \leq \theta_i(A).$$

3 A new existence condition

In this section, we will give a new existence condition, Theorem 4, for distance-regular graphs. To do this we first show Lemma 2 and Proposition 3.

Lemma 2 Let Γ be a distance-regular graph with valency k and diameter $D \geq 2$. Let x be a vertex of Γ and let \bar{C} be a co-clique of size $s \geq 2$ in the local graph $\Delta(x)$ at x . Then

$$c_2 - 1 \geq \frac{s(a_1 + 1) - k}{\binom{s}{2}}.$$

Proof: Let $V(\bar{C}) = \{y_1, y_2, \dots, y_s\}$. Since $d_\Gamma(y_i, y_j) = 2$, $|\Gamma(x, y_i, y_j)| \leq c_2 - 1$ holds for any $i \neq j$. Then by the principle of inclusion and exclusion,

$$\begin{aligned} k = |\Gamma(x)| &\geq \left| \bigcup_{i=1}^s (\Gamma(x, y_i) \cup \{y_i\}) \right| \\ &\geq \sum_{i=1}^s |\Gamma(x, y_i) \cup \{y_i\}| - \sum_{1 \leq i < j \leq s} |\Gamma(x, y_i, y_j)| \\ &\geq s(a_1 + 1) - \binom{s}{2}(c_2 - 1). \end{aligned}$$

□

Proposition 3 Let Γ be a distance-regular graph with valency k and diameter $D \geq 2$. Let s be maximal such that for all x and all $y, z \in \Gamma(x)$ with $y \approx z$, there exists a co-clique of size at least s in $\Delta(x)$ containing y and z . Then

$$(i) \quad s \geq \frac{k}{a_1 + 1}$$

$$(ii) \quad c_2 - 1 \geq \max\left\{\frac{s'(a_1 + 1) - k}{\binom{s'}{2}} \mid 2 \leq s' \leq s\right\} \text{ and equality implies } \Gamma \text{ is a Terwilliger graph.}$$

Proof: Let s be maximal satisfying the condition in the Proposition 3. Then $k \leq s(a_1 + 1)$ as $\Delta(x)$ has valency a_1 and k vertices. This shows (i). By Lemma 2, the inequality in (ii) holds. Next, we assume that the equality holds in (ii). Let

$2 \leq s'' \leq s$ be an integer satisfying $\frac{s''(a_1+1)-k}{\binom{s''}{2}} = \max\{\frac{s'(a_1+1)-k}{\binom{s'}{2}} \mid 2 \leq s' \leq s\}$, then by Lemma 2 there exists a co-clique \bar{C}'' on $\{y_1, y_2, \dots, y_{s''}\}$ such that for any two vertices y_i, y_j at distance two, $|\Gamma(x, y_i, y_j)| = c_2 - 1$ holds. That is, if we take three vertices z_1, z_2 and z_3 such that $d_\Gamma(z_2, z_3) = 2$, $z_1 \sim z_2$ and $z_1 \sim z_3$, then since $z_1 \in \Gamma(z_2, z_3)$ and $|\Gamma(z_1, z_2, z_3)| = c_2 - 1$, the valency of z_1 in $\Gamma(z_2, z_3)$ is $c_2 - 1$. Hence the subgraph induced by $\Gamma(z, w)$ is a clique of size c_2 for any two vertices z and w at distance two in Γ . So, Γ is a Terwilliger graph. \square

For all known examples of Terwilliger graphs, we have equality in case (ii) above.

Theorem 4 Let Γ be a distance-regular graph with valency k and diameter $D \geq 2$ and define $\alpha = \lceil \frac{k}{a_1+1} \rceil$. Then $c_2 - 1 \geq \frac{\alpha(a_1+1)-k}{\binom{\alpha}{2}}$ and equality implies that Γ is a Terwilliger graph.

Proof: This is an immediate consequence of Proposition 3. \square

Theorem 4 gives the new existence condition for distance-regular graphs and the following two intersection arrays in [1, p.425-431] are ruled out.

Corollary 5 There are no distance-regular graphs with one of the following intersection arrays:

$$(i) \{44, 30, 5; 1, 3, 40\}; \quad (ii) \{65, 44, 11; 1, 4, 55\}.$$

Proof:

- (1) Since a distance-regular graph Γ with intersection array (i) satisfies $c_2 - 1 = 2 = \frac{\alpha(a_1+1)-k}{\binom{\alpha}{2}}$, where $\alpha = \lceil \frac{k}{a_1+1} \rceil = 4$, Γ is a Terwilliger graph by Theorem 4. But this is impossible by [1, Corollary 1.16.6].
- (2) As $\lceil \frac{65}{20+1} \rceil = 4$, there is no distance-regular graph with intersection array (ii), by Theorem 4.

\square

Remark: A. Jurišić and J. Koolen [4] proved that distance-regular graphs with intersection arrays $\{81, 56, 24, 1; 1, 3, 56, 81\}$, $\{117, 80, 30, 1; 1, 6, 80, 117\}$, $\{117, 80, 32, 1; 1, 4, 80, 117\}$, and $\{189, 128, 45, 1; 1, 9, 128, 189\}$ do not exist. It also follows from Theorem 4.

4 Shilla distance-regular graphs

In this section we first give a lower bound on the second largest eigenvalue of a distance-regular graph with diameter 3. Then we will define Shilla distance-regular graphs and give some results on them.

Lemma 6 Let Γ be a distance-regular graph with valency k and diameter 3. Then the second largest eigenvalue θ_1 of Γ satisfies

$$\theta_1 \geq \min\left\{\frac{a_1 + \sqrt{a_1^2 + 4k}}{2}, a_3\right\}.$$

Proof: Let x be a vertex of Γ . As the induced subgraph on $\{x\} \cup \Gamma(x)$ (respectively $\Gamma_3(x)$) has largest eigenvalue $\frac{a_1 + \sqrt{a_1^2 + 4k}}{2}$ (respectively a_3), it follows that the induced subgraph on $\{x\} \cup \Gamma(x) \cup \Gamma_3(x)$ has the second largest eigenvalue at least $\min\left\{\frac{a_1 + \sqrt{a_1^2 + 4k}}{2}, a_3\right\}$. Now the lemma follows by Theorem 1. \square

Theorem 7 Let θ be an eigenvalue of a distance-regular graph Γ with valency k and diameter 3. Then the following are equivalent:

(i) $u_2(\theta) = 0$;

(ii) $\theta = a_3$;

(iii) $\theta = \frac{a_1 + \sqrt{a_1^2 + 4k}}{2}$.

Moreover, if (i) – (iii) hold then θ is the second largest eigenvalue of Γ .

Proof: (i) \Leftrightarrow (ii): Let $u_0 = 1$, u_1 , u_2 , u_3 be the standard sequence for θ . As $c_3u_2 + a_3u_3 = \theta u_3$, it follows $u_2 = 0$ if and only if $\theta = a_3$.

(i) \Leftrightarrow (iii): As $1 + a_1u_1 + b_1u_2 = \theta u_1$ and $u_1 = \frac{\theta}{k}$, it follows that $\theta = \frac{a_1 + \sqrt{a_1^2 + 4k}}{2}$ if and only if $u_2 = 0$.

Moreover, if (i) – (iii) hold then, $u_0 = 1 > 0$, $u_1 = \frac{a_3}{k} > 0$, $u_2 = 0$, and $u_3 = -\frac{c_2a_3}{kb_2} < 0$ and hence θ is the second largest eigenvalue of Γ by [1, Corollary 4.1.2]. \square

A distance-regular graph with diameter 3 and valency k is called *Shilla* if its second largest eigenvalue θ_1 satisfies $\theta_1 = a_3$. It follows by Theorem 7 that $\theta_1 = a_3 = \frac{a_1 + \sqrt{a_1^2 + 4k}}{2}$ and hence $k = (a_3 - a_1)a_3$. For a Shilla distance-regular graph Γ , put $b = b(\Gamma) := a_3 - a_1$. Then clearly $b \geq 2$ and $k = ba_3$.

A Shilla distance-regular graph Γ with $b(\Gamma) = b$ has distinct four eigenvalues $\theta_0 = k = ba_3 > \theta_1 = a_3 > \theta_2 > \theta_3$, where θ_2 and θ_3 are two roots of the equation

$x^2 - (a_1 + a_2 - k)x + (b - 1)b_2 - a_2 = 0$. Let m_i be the multiplicity of θ_i . If both θ_2 and θ_3 are integers then $(a_1 + a_2 - k)^2 - 4((b - 1)b_2 - a_2)$ is a perfect square. If both θ_2 and θ_3 are non-integers, then $m_2 = m_3$ holds. This implies, by Equation 1, that the equation

$$(b_2 + c_2)(b_2 + c_2 - a_3)(b_2 + c_2 + (b - 1)a_3) - bb_2^2 + (2b - 3)c_2^2 + b(b - 1)c_2 + (b - 1)^2 a_3 c_2 - b(b - 1)a_3 b_2 + (b - 3)b_2 c_2 = 0 \quad (2)$$

holds. In Theorem 11 below, we will discuss the situation $m_2 = m_3$ in more detail.

Now, we will show that there are finitely many Shilla distance-regular graphs Γ with fixed $b(\Gamma)$. To do this we first show Lemma 8.

Lemma 8 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = b$. Then,

$$c_2 \geq \frac{2a_3 - b^2 + b + 2}{b(b + 1)}.$$

Proof: Let x be a vertex of Γ . Then there exists a co-clique of size $b + 1$ in $\Delta(x)$ as $k = ba_3 = b(a_1 + b) > b(a_1 + 1)$ and by Lemma 2, the proof is complete. \square

Theorem 9 For given $\beta \geq 2$, there are finitely many Shilla distance-regular graphs Γ with $b(\Gamma) = \beta$.

Proof: For given $\beta \geq 2$, let Γ be a Shilla distance-regular graph with valency k , $b(\Gamma) = \beta$ and n vertices. Then clearly $k = \beta a_3 = \beta(a_1 + \beta) = \beta(a_1 + 1) + \beta^2 - \beta$. We will show that k is bounded above by β . We first show the following.

Claim: $k < \beta^3 - \beta$ or $n < k(2\beta^3 - \beta + 1)$.

Proof of claim: If $a_1 + 1 < \beta^2 - \beta$, then $k = \beta a_3 = \beta(a_1 + \beta) < \beta^3 - \beta$. So, let us assume $a_1 + 1 \geq \beta^2 - \beta$ then clearly $k \geq \beta^3 - \beta$. Lemma 8 implies $c_2 \geq \frac{a_3 + (a_3 + 1 - \beta^2) + \beta + 1}{\beta(\beta + 1)} > \frac{a_3 + 1}{\beta(\beta + 1)}$, where the second inequality follows from $a_3 + 1 \geq \beta^2$. As $c_3 = (\beta - 1)a_3$ and $b_1 = (\beta - 1)(a_3 + 1)$, it follows that

$$\begin{aligned} n &= 1 + k + k \frac{b_1}{c_2} + k \frac{b_1 b_2}{c_2 c_3} = 1 + k + k \frac{(\beta - 1)(a_3 + 1)}{c_2} + k \frac{b_2}{c_2} + \frac{\beta b_2}{c_2} \\ &\leq 1 + k + 2k\beta(\beta - 1)(\beta + 1) + \beta^2(\beta^2 - 1) \leq 1 + k + 2k\beta(\beta - 1)(\beta + 1) + k\beta \\ &< k(2\beta^3 - \beta + 2). \end{aligned}$$

So, the claim is proved.

Now by letting m_1 to be the multiplicity of $\theta_1 = a_3$, it follows from Equation 1 that $m_1 < \frac{n}{u_1(a_3)^2 k}$. By [1, Theorem 5.3.2], $\sqrt{k} < m_1$ holds. As $m_1 < \frac{n}{u_1(a_3)^2 k}$

$< \frac{k(2\beta^3 - \beta + 2)}{u_1(a_3)^2 k} = 2\beta^5 - \beta^3 + 2\beta^2$, it follows $k < 4\beta^{10}$. This shows the theorem. \square

In the next result, we give some divisibility conditions for Shilla distance-regular graphs.

Lemma 10 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = b$. Then the following holds:

- (i) c_2 divides $(b-1)a_3b_2$;
- (ii) c_2 divides $(b-1)ba_3(a_3+1)$;
- (iii) c_2 divides $b(a_3+1)b_2$;
- (iv) c_2 divides $(b+a_3)b_2$ and $(b+a_3)b_2 \geq (1+a_3)c_2$, where equality is attained if and only if $p_{33}^3 = 0$;
- (v) c_2 divides $(b-1)bb_2$.

Proof: (i) follows from the fact that p_{32}^3 is a non-negative integral.

(ii) and (iii) hold as $k_2 = \frac{kb_1}{c_2}$ and $k_3 = \frac{kb_1b_2}{c_2c_3}$ are integral respectively. Since $\frac{kb_1b_2}{c_2c_3} = k_3 = p_{30}^3 + p_{31}^3 + p_{32}^3 + p_{33}^3 = 1 + a_3 + \frac{c_3(b_2-1)+a_3(a_3-1-a_1)}{c_2} + p_{33}^3$ is integral, it follows that $\frac{(b+a_3)b_2}{c_2} \geq 1 + a_3$ and equality is attained if and only if $p_{33}^3 = 0$. Since $\frac{(b+a_3)b_2}{c_2}$ and $\frac{(b-1)(b+a_3)b_2}{c_2}$ are integral, (v) holds by (i). \square

We will give some necessary conditions for Shilla distance-regular graphs with $m_2 = m_3$.

Theorem 11 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = b \geq 2$ and $m_2 = m_3$. Then the following holds:

- (i) $a_3 - b < b_2 + c_2 < a_3 + b$;
- (ii) $c_2 < b_2 + b$;
- (iii) If $b_2 + c_2 = a_3$ or $b_2 = c_2$, then $a_3 = \frac{b(b-1)}{2}$.

Proof: Since $m_2 = m_3$, Equation(2) holds.

(i): If $b_2 + c_2 \leq a_3 - b$, then the LHS of Equation(2) is negative. Hence $a_3 - b < b_2 + c_2$. In similar fashion, we can show $b_2 + c_2 < a_3 + b$. Thus, $a_3 - b < b_2 + c_2 < a_3 + b$.

(ii): If $c_2 \geq b_2 + b$, then Lemma 10 (iv) implies $(b+a_3)b_2 \geq (1+a_3)c_2 \geq (1+a_3)(b_2+b)$. This means, $b_2 \geq \frac{b}{b-1}(a_3+1) = a_3+1 + \frac{a_3+1}{b-1} > a_3+2$, where the last inequality holds by $a_3 \geq b$. So, $b_2 + c_2 > a_3 + b$ and this is a contradiction to (i). Thus $c_2 < b_2 + b$.

(iii) If $b_2 + c_2 = a_3$, then Equation(2) becomes $b^2(b_2^2 - c_2^2) + 2c_2(b_2 + c_2) = b(b-1)c_2$ and it follows $b_2 \leq c_2$. If $b_2 = c_2$, then $c_2 = \frac{b(b-1)}{4}$ and hence $a_3 = \frac{b(b-1)}{2}$. If $b_2 < c_2$, then

$b(b-1)c_2 = b^2(b_2^2 - c_2^2) + 2c_2(b_2 + c_2) \leq 4c_2^2 - 2c_2 - 2b^2c_2 + b^2$ and hence $c_2 \geq \frac{2b^2-b+2}{4}$. Now it follows from Lemma 10 that $\frac{(b+a_3)b_2}{c_2} = \frac{(b+b_2+c_2)b_2}{c_2}$ is integral and hence $\frac{(b_2+b)b_2}{c_2}$ is integral. Since $b_2 = c_2 - \alpha$ for some $1 \leq \alpha < b$, we find c_2 divides $\alpha(b - \alpha)$. Hence $c_2 \leq \alpha(b - \alpha) \leq \frac{b^2}{4}$, but this contradicts $c_2 \geq \frac{2b^2-b+2}{4}$. If $b_2 = c_2$, then Equation(2) becomes $2c_2(2c_2 - a_3)(2c_2 + (b-1)a_3) + b(b-1)c_2 - (b-1)a_3c_2 + 2(b-3)c_2^2$ and this is always positive (respectively negative) if $2c_2 > a_3$ (respectively $2c_2 < a_3$). Thus $2c_2 = a_3$ and hence $a_3 = \frac{b(b-1)}{2}$. \square

Note that in case (iii) above, we have the intersection arrays

$$\left\{ \frac{b^2(b-1)}{2}, \frac{(b-1)(b^2-b+2)}{2}, \frac{b(b-1)}{4}; 1, \frac{b(b-1)}{4}, \frac{b(b-1)^2}{2} \right\}$$

and they are only feasible for $b \equiv 0, 1 \pmod{4}$. Besides this family of intersection arrays, using the computer, the only other feasible intersection arrays for Shilla distance-regular graphs with $m_2 = m_3$ and $a_3 \leq 100$ are:

- (*) (i) $\{120, 117, 20; 1, 1, 108\}$; (ii) $\{676, 675, 31; 1, 9, 650\}$;
 (iii) $\{486, 440, 50; 1, 10, 432\}$; (iv) $\{4264, 4233, 102; 1, 17, 4182\}$.

In the next theorem, we classify Shilla distance-regular graphs Γ with $b(\Gamma) = 2$.

Theorem 12 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = 2$. Then Γ is one of the following graphs:

- (i) the Odd graph with valency 4;
- (ii) a generalized hexagon of order (2,2);
- (iii) the Hamming graph $H(3, 3)$;
- (iv) the Doro graph with intersection array $\{10, 6, 4; 1, 2, 5\}$;
- (v) the Johnson graph $J(9, 3)$.

Proof: If $b = 2$, then $b_1 = a_3 + 1$ and hence $\theta_1 = a_3 = b_1 - 1$. By [1, Theorem 4.4.11] we only need to consider the cases $c_2 = 1$ or Γ is one of a Doob, a Hamming, a locally Petersen, a Johnson, a halved cube, or the Gosset graph. By [1, Theorem 1.16.5] the only locally Petersen graph which is a Shilla distance-regular graph is the Doro graph. If Γ is a Doob or a Hamming graph, then $c_3 = 3 = a_3$ and it follows that $\Gamma = H(3, 3)$. If Γ is a Johnson graph, then $c_3 = 9 = a_3$ and it follows that $\Gamma = J(9, 3)$. Neither the Gosset graph nor the halved 6-cube are possible as they have $a_3 = 0$. Also, the halved 7-cube is not a Shilla distance-regular graph. To complete the proof of this theorem, we only need to consider the case $c_2 = 1$. If $c_2 = 1$, then Γ is a locally disjoint union of $(a_1 + 1)$ -cliques. This implies that $a_1 + 1$ divides k , and hence $a_3 \in \{2, 3\}$. If $a_3 = 2$, then $k = 4, c_1 = 1, a_1 = 0, b_1 = 3, c_3 = 2$ and $b_2 \in \{1, 2, 3\}$. Only for $b_2 = 3$, the multiplicity m_1 is an integer, and Γ is the Odd graph with valency 4. If $a_3 = 3$, then

$k = 6, c_1 = 1, a_1 = 1, b_1 = 4$, and $b_2 \in \{1, 2, 3, 4, 5\}$. Only for $b_2 = 4$, the multiplicity m_1 is an integer, and Γ is a generalized hexagon of order $(2, 2)$. \square

The following lemma gives a sufficient condition for a distance-regular graph to be Q -polynomial.

Lemma 13 Let Γ be a distance-regular graph with diameter $D = 3$, n vertices and eigenvalues $k > \theta_1 > \theta_2 > \theta_3$. If $n \geq \frac{(m_1+2)(m_1+1)}{2}$, then $q_{11}^2 = 0$ or $q_{11}^3 = 0$ and hence Γ is Q -polynomial with respect to θ_1 , where q_{11}^2 and q_{11}^3 are Krein parameters.

Proof: As $n \geq \frac{(m_1+2)(m_1+1)}{2}$ and $n = 1 + m_1 + m_2 + m_3$, it follows that $m_2 + m_3 \geq \binom{m_1+1}{2}$. As $\sum_{q_{11}^i \neq 0} m_i \leq \binom{m_1+1}{2}$ [1, Proposition 4.1.5] and $q_{11}^0 > 0$, it follows that $q_{11}^2 = 0$ or $q_{11}^3 = 0$. This implies that Γ is Q -polynomial with respect to θ_1 . \square

Lemma 14 Let Γ be a Shilla distance-regular graph with n vertices, $b(\Gamma) = \beta$ and valency k . If Γ is not Q -polynomial with respect to θ_1 then $k < \beta^5(\beta + 1)^2$.

Proof: Let $k \geq \beta^3 - \beta$. If Γ is not Q -polynomial with respect to θ_1 then $n < \frac{(m_1+1)(m_1+2)}{2}$ by Lemma 13. By Equation 1 we find $m_1 + 2 < \frac{n}{a_3/\beta} + 2 < \frac{\beta+1}{\beta} \frac{n}{a_3/\beta}$, where the last inequality holds by $\frac{n}{a_3} > \frac{k}{a_3} = \beta \geq 2$. Combining the above two inequalities we find $\sqrt{2n} < \frac{\beta+1}{\beta} \frac{n}{a_3/\beta}$ and hence $2(\frac{a_3}{\beta+1})^2 < n$. As $k \geq \beta^3 - \beta$, by Claim in Theorem 9, we find $n < k(2\beta^3 - \beta + 1)$. So $2(\frac{a_3}{\beta+1})^2 < k(2\beta^3 - \beta + 1)$. Since $k = \beta a_3$, we find $k < \beta^5(\beta + 1)^2$. \square

Proposition 15 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = b$. Then the following holds:

- (i) $q_{11}^2 > 0$;
- (ii) $q_{11}^3 \geq 0$ if and only if $\theta_3 \geq -\frac{b(bb_2+c_2)}{b_2+c_2}$.

So, in particular $\theta_3 \geq -\frac{b(bb_2+c_2)}{b_2+c_2}$ holds.

Proof: Note that $q_{jh}^i \geq 0 = \frac{m_j m_h}{|V\Gamma|} \sum_{l=0}^D k_l u_l(\theta_i) u_l(\theta_j) u_l(\theta_h) \geq 0$ [1, Proposition 4.1.5].

Hence $q_{11}^i \geq 0$ if and only if $\sum_{l=0}^3 k_l u_l(\theta_1)^2 u_l(\theta_i) \geq 0$ if and only if

$$c_2 \theta_i^3 - c_2(a_1 + a_2) \theta_i^2 + (c_2 a_1 a_2 - b_1 c_2^2 - c_2 k b_2^2 c_3) \theta_i + c_2 a_2 k + b^2 b_2^2 c_3 \geq 0$$

Since θ_2 and θ_3 are two roots of polynomial $\theta^2 - (a_1 + a_2 - k)\theta + (b-1)b_2 - a_2$, we obtain that $q_{11}^i \geq 0$ if and only if $(b^2b_2 + bc_2 + b_2\theta_i + c_2\theta_i) \geq 0$ for $i = 2, 3$. As, $\theta_2 > \theta_3$ we see immediately that $q_{11}^2 > 0$ and also that $q_{11}^3 \geq 0$ if and only if $\theta_3 \geq -\frac{b(bb_2+c_2)}{b_2+c_2}$. This shows the proposition. \square

Corollary 16 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = b$. Then

$$-b^2 < \theta_3 < -b$$

Proof: Let x be a vertex in Γ . Then the induced subgraph on $\{x\} \cup \Gamma(x)$ has two eigenvalues, $-b$ and a_3 . Then, by Theorem 1, $\theta_3 \leq -b$ holds. But if $\theta_3 = -b$, then $u_2(\theta_3) = 0$ and this is not possible by Theorem 7. The lower bound follows immediately from Proposition 15. \square

We will improve the lower bound for θ_3 in Theorem 20 below.

Corollary 17 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = b$. Then Γ is Q -polynomial with respect to θ_1 if and only if $\theta_3 = -\frac{b(bb_2+c_2)}{b_2+c_2}$. If Γ is Q -polynomial with respect to θ_1 , then all eigenvalues of Γ are integral, $b_2 + c_2$ divides $b(b-1)b_2$ and $-b^2 + 1 \leq \theta_3 \leq -\frac{b^2(b+3)}{3b+1}$.

Proof: Note that Γ is Q -polynomial with respect to θ_1 if and only if $q_{11}^2 = 0$ or $q_{11}^3 = 0$. Hence the first part of this corollary follows from Proposition 15. Assume Γ is Q -polynomial with respect to θ_1 . Then $\theta_3 = -b - \frac{b(b-1)b_2}{b_2+c_2}$ and hence θ_3 is integral. So, all the eigenvalues are integral. As $b \leq a_3$, it follows that $c_2 \leq \frac{(b+a_3)b_2}{1+a_3} \leq \frac{2bb_2}{1+b}$ by Lemma 10. So, $\theta_3 \leq -b - \frac{b(b-1)(b+1)}{3b+1} = -\frac{b^2(b+3)}{3b+1}$. Thus $-b^2 + 1 \leq \theta_3 \leq -\frac{b^2(b+3)}{3b+1}$ holds by Corollary 16. \square

In the next two results, we classify the Shilla distance-regular graphs Γ with $b(\Gamma) = 3$.

Proposition 18 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = 3$ and let Γ be Q -polynomial with respect to θ_1 . Then Γ has one of the following intersection arrays.

$$(i) \quad \{42, 30, 12; 1, 6, 28\}, \quad (ii) \quad \{105, 72, 24; 1, 12, 70\}.$$

Proof: By Corollary 17, if Γ is a Q -polynomial with respect to θ_1 , then $\theta_3 \in \{-6, -7, -8\}$.

If $\theta_3 = -6$, then $c_2 = b_2$. Since θ_3 is a root of the equation $x^2 - (a_1 + a_2 - k)x + (b-1)b_2 - a_2 = 0$, $b_2 = c_2$ and $b(\Gamma) = b = 3$, it follows $a_3 = \frac{8}{3}b_2 - 6$. But this means $m_1 = 33 - \frac{135}{2b_2}$ what is impossible, as m_1 must be an integer.

If $\theta_3 = -7$, Similarly, we obtain $b_2 = 2c_2$ and $a_3 = \frac{7}{2}c_2 - 7$. Then $m_1 = 60 - \frac{108}{c_2}$. Since a_3 and m_1 have to be integers, it follows that 2 divides c_2 and c_2 divides 108, that is, $c_2 \in \{2, 4, 6, 12, 18, 36, 54, 108\}$. The case $c_2 = 2$ implies $a_3 = 0$, which is impossible. For $c_2 \in \{18, 36, 54, 108\}$, we find that m_3 is non-integral. The case $c_2 = 4$ gives us the intersection array $\{21, 16, 8; 1, 4, 14\}$ and it was shown by K. Coolsaet[2] that a distance-regular graph with this intersection array does not exist. The cases $c_2 = 6$ and $c_2 = 12$ give the intersection arrays (i) and (ii) respectively.

If $\theta_3 = -8$, Similarly, we obtain $b_2 = 5c_2$ and $a_3 = \frac{32}{5}c_2 - 8$. But this means $m_1 = 141 - \frac{315}{2c_2}$ what is impossible.

□

Theorem 19 Let Γ be a Shilla distance-regular graph with $b(\Gamma) = 3$. Then Γ has one of the following intersection arrays.

- | | | |
|-----------------------------------|-------------------------------------|--------------------------------------|
| (i) $\{12, 10, 5; 1, 1, 8\}$, | (ii) $\{12, 10, 2; 1, 2, 8\}$, | (iii) $\{12, 10, 3; 1, 3, 8\}$, |
| (iv) $\{15, 12, 6; 1, 2, 10\}$, | (v) $\{24, 18, 9; 1, 1, 16\}$, | (vi) $\{27, 20, 10; 1, 2, 18\}$, |
| (vii) $\{30, 22, 9; 1, 3, 20\}$, | (viii) $\{42, 30, 12; 1, 6, 28\}$, | (ix) $\{60, 42, 18; 1, 6, 40\}$, |
| (x) $\{69, 48, 24; 1, 4, 46\}$, | (xi) $\{93, 64, 24; 1, 6, 62\}$, | (xii) $\{105, 72, 24; 1, 12, 70\}$. |

Note that all the above intersection arrays have $\theta_3 \geq -7$

Proof: If Γ is Q -polynomial with respect to θ_1 then it follows from Proposition 18 that Γ has intersection array (viii) or (xii). If Γ is not Q -polynomial with respect to θ_1 then by Lemma 14, $a_3 < 3^4 \times 4^2 = 1296$. We checked by computer that the above arrays are the only possible intersection arrays for Shilla distance-regular graphs with $a_3 < 1296$. □

Remark: The unitary nonisotropics graph with $q = 4$ as defined in [1, Section 12.4] has intersection array (i). It is not known whether it is unique or not. There exists a unique distance-regular graph with intersection array (iii) namely the Doro graph as defined in [1, Section 12.1]. For the other intersection arrays, it is not known whether a distance-regular graph with those intersection arrays does exist, or not.

Now, we improve the lower bound of the smallest eigenvalue θ_3 for a Shilla distance-regular graph.

Theorem 20 For a Shilla distance-regular graph with $b(\Gamma) = b$ and smallest eigenvalue θ_3 , we have $\theta_3 < -b^2 + 2$ if and only if $b = 2$.

Proof: (\Leftarrow) For $b = 2$, we are done by Theorem 12.

(\Rightarrow) Let $\theta_3 < -b^2 + 2$. Then by Theorems 12 and 19 we have $b = 2$ or $b \geq 4$. So, let us assume $b \geq 4$. By Corollary 16, we have $-b^2 < \theta_3 < -b^2 + 2$. Then either $\theta_3 = -b^2 + 1$ or $m_2 = m_3$ and θ_3 is non-integer. If $\theta_3 = -b^2 + 1$, then by Proposition 15, $b_2 \geq (b^2 - b - 1)c_2$. Since $\theta_3 = -b^2 + 1$ is a root of the equation $x^2 - (a_1 + a_2 - k)x + (b - 1)b_2 - a_2 = 0$ and $b_2 \geq (b^2 - b - 1)c_2$, we have

$$a_3 = b_2 + \frac{b^2 - 2}{b^2 - b - 1}c_2 - (b^2 - 1) \geq \frac{(b-1)^3(b+1)}{(b^2 - b - 1)}c_2 - (b^2 - 1)$$

and hence $c_2 \in \{1, 2, 3\}$ by Lemma 10. Since $a_3 = b_2 + \frac{b^2 - 2}{b^2 - b - 1}c_2 - (b^2 - 1) \in \mathbf{Z}$, $b^2 - b - 1$ divides $(b - 1)c_2$ and hence $b^2 - b - 1 \leq 3(b - 1)$. This contradicts $b \geq 4$. Let us now consider the case $m_2 = m_3$ and θ_3 is non-integer. As $c_2 + b_2 \leq a_3 + b - 1$, by Theorem 11, the LHS of Equation (2) is at most

$$(**) \quad (b_2 + c_2)((3b - 4)c_2 - b_2) + (b - 1)((2b - 2)c_2 - b_2)a_3 + b(b - 1)c_2.$$

Since $b_2 \geq (3b - 4)c_2$ implies that $(**)$ is negative, we have $b_2 < (3b - 4)c_2$. By Proposition 15, we have $\theta_3 \geq -b^2 + \frac{b(b-1)c_2}{b_2 + c_2} \geq -b^2 + \frac{b}{3}$, and hence $4 \leq b \leq 5$. Let us consider first $b = 5$. As $-23 > \theta_3 \geq -25 + \frac{20c_2}{b_2 + c_2}$, it follows that $9c_2 < b_2 < 11c_2$. As $(**) > 0$, it follows that $a_3 \leq 13$. As the intersection array $\{50, 44, 5; 1, 5, 40\}$ has $\theta_3 = -7.623$, this case follows now from $(*)$. Secondly, we assume that $b = 4$. As $-14 > \theta_3 \geq -16 + \frac{12c_2}{b_2 + c_2}$, it follows that $5c_2 < b_2 < 8c_2$. Since $b_2 + c_2 \leq a_3$ implies that the LHS of Equation (2) is negative, we have $a_3 < b_2 + c_2 \leq a_3 + 3$. If $b_2 + c_2 = a_3 + 1$ then the LHS of Equation (2) $= 4a_3^2 + 5a_3 + 1 + 9a_3c_2 - 12a_3b_2 - 4b_2^2 + 5c_2^2 + 12c_2 + b_2c_2$. Since $5c_2 \leq b_2$, $b_2^2 \geq 5c_2^2 + 12c_2 + b_2c_2$ and $6b_2 \geq 5a_3$. Hence the LHS of Equation (2) is negative. Similarly, for $b_2 + c_2 = a_3 + 2$, the LHS of Equation (2) is negative if $a_3 \geq 9$. So, $a_3 \leq 8$ and then we are done by $(*)$. If $b_2 + c_2 = a_3 + 3$ then the LHS of Equation (2) $= 2(13c_2 - 2b_2)(c_2 + b_2) + 3(3b_2 - 5c_2 - 9)$. Since $b_2 \leq \frac{19}{3}c_2$ implies that the LHS of Equation (2) is positive whenever $b_2 + c_2 \geq 45$ (i.e. $a_3 \geq 42$), either $(b_2 \leq \frac{19}{3}c_2$ and $a_3 \leq 41)$ or $b_2 > \frac{19}{3}c_2$. By $(*)$ we obtain $b_2 > \frac{19}{3}c_2$. Since $b_2 \geq \frac{27}{4}c_2$ implies that the LHS of Equation (2) is negative whenever $c_2 \geq 9$, either $(b_2 \geq \frac{27}{4}c_2$ and $c_2 \leq 8)$ or $b_2 < \frac{27}{4}c_2$. As $c_2 \leq 8$ implies that $a_3 \leq 85$ by Lemma 8, from $(*)$ we obtain $\frac{19}{3}c_2 < b_2 < \frac{27}{4}c_2$. Then by Lemma 10 (v), $\frac{12b_2}{c_2} \in \{77, 78, 79, 80\}$. In the first two possibilities the LHS of Equation (2) is negative. In the last two possibilities the number c_2 is non-integral. This shows the theorem. \square

Acknowledgments

Section 3 was inspired by a conversation with Sejeong Bang. We also would like to thank her for the careful reading she did. Comments by Mitsugu Hirasaka are appreciated much. This work is partially supported by KRF-2008-314-C00007 and this support is greatly appreciated.

References

- [1] A.E. Brouwer, A.M. Cohen and A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, 1989.
- [2] K. Coolsaet, A distance-regular graph with intersection array $\{21, 16, 8; 1, 4, 14\}$ does not exist, European J. Combin. 26 (2005), 709-716 .
- [3] W.H. Haemers, Interlacing eigenvalues and graphs, Linear Algebra Appl. 226/228 (1995), 593-616.
- [4] A. Jurišić and J. Koolen, Nonexistence of some antipodal distance-regular graphs of diameter four, European J. Combin. 21 (2000), 1039-1046.